

Multiple Lambda cosmology: dark fluid with time-dependent equation of state as classical analog of cosmological landscape

Shin'ichi Nojiri^{1,*} and Sergei D. Odintsov^{†2,‡}

¹*Department of Physics, Nagoya University, Nagoya 464-8602, Japan*

²*Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Ciències de l'Espai (IEEC-CSIC), Campus UAB, Facultat de Ciències, Torre C5-Par-2a pl, E-08193 Bellaterra (Barcelona), Spain*

We discuss FRW universe with time-dependent EoS dark fluid which leads to multiple de Sitter space. This model (as well as its scalar-tensor version) may be considered as some classical analog of cosmological landscape. The universe expansion history may look as transitions between different deSitter eras which suggests the interesting solution for cosmological constant problem. For specific time-dependent EoS dark fluid the possibility of transitions between universe regions with positive and negative cosmological constant is also established.

PACS numbers: 11.25.-w, 95.36.+x, 98.80.-k

1. In the study of celebrated dark energy problem (for a review, see [1]) the ideal fluid with specific (sometimes, strange) equation of state (EoS) remains to be the simplest possibility for the description of the current cosmic acceleration. Various examples of ideal fluid EoS may be considered for this purpose: constant EoS with negative pressure, imperfect EoS [2], general EoS [3], inhomogeneous EoS [4] (with time-dependent bulk viscosity as particular case [5]) and so on (for extensive review, see [1]). Trying to save General Relativity and to explain the cosmic acceleration at the same time may lead to the conjecture about even more exotic dark fluids. Moreover, General Relativity with ideal fluid of any type may be rewritten in the equivalent form [6] as modified gravity (for general review, see [7]). Of course, the introduction of ideal fluid with complicated equation of state is somehow phenomenological approach because no explanation of the origin of such dark fluid follows.

The interesting possibility to explain the dark fluid origin may be related with string theory (for recent review of some attempts in this direction, see [8]). The following sequence may be proposed: string/M-theory is approximated by modified (super)gravity [7] which is observed finally as General Relativity with exotic dark fluid. If such conjecture is (even partially) true, it is expected that some string-related phenomena may be typical in dark energy universe. One celebrated stringy effect possibly related with early universe is string landscape (see, for instance, [9]) which may lead to some observational consequences (see, for instance, [10]). The cosmological landscape may be responsible for discrete mass spectrum of scalar/spinor equations [11].

In the present Letter we suggest the dark fluid with time-dependent EoS which may be considered as simple classical analog of string landscape. It leads to multiple de Sitter space at the same moment. It may also describe transitions between de Sitter universes, suggesting the way to solve the cosmological constant problem. Unlike to proper string landscape the true ground state may be still selected because the on-shell action is not the same for all these de Sitter solutions. Some modification of dark fluid EoS suggests the possible transition between regions with positive and negative cosmological constant.

2. Let us study the specific model of ideal fluid which leads to multiple Lambda cosmology. The FRW equations are given by

$$\frac{3}{\kappa^2}H^2 = \rho, \quad -\frac{2}{\kappa^2}\dot{H} = p + \rho. \quad (1)$$

Here ρ is the energy density and p is the pressure. Instead of (1), one may include the cosmological constant

[†] also at Lab. Fundam. Study, Tomsk State Pedagogical University, Tomsk

*Electronic address: nojiri@phys.nagoya-u.ac.jp

‡Electronic address: odintsov@ieec.uab.es

from gravity:

$$\frac{3}{\kappa^2}H^2 = \rho + \frac{\Lambda}{\kappa^2}, \quad -\frac{2}{\kappa^2}\dot{H} = p + \rho. \quad (2)$$

We can, however, redefine ρ and p to absorb the contribution from the cosmological constant:

$$\rho \rightarrow \rho - \frac{\Lambda}{\kappa^2}, \quad p \rightarrow p + \frac{\Lambda}{\kappa^2}. \quad (3)$$

By the redefinition (3), we reobtain (1). Hence, it is enough to consider only dark fluid in FRW equation.

If ρ and p are given in terms of the function $f(q)$ with a parameter q as (compare with similar Anzats [12])

$$\rho = \frac{3}{\kappa^2}f(q)^2, \quad p = -\frac{3}{\kappa^2}f(q)^2 - \frac{2}{\kappa^2}f'(q), \quad (4)$$

the following solution of Eq.(1) exists:

$$H = f(t). \quad (5)$$

Note that the origin of the time can be chosen arbitrary. In (5), $t = q$ but one may choose $t = q + t_0$ with an arbitrary constant t_0 . This shows that besides the solution (5), $H = f(t - t_0)$ can be a solution.

If we delete q in (4), we have a general equation of state (EoS):

$$F(\rho, p) = 0. \quad (6)$$

In case that $f'(q) = 0$ has a solution $q = q_0$, there is a solution where H is a constant:

$$H = H_0 \equiv f(q_0), \quad (7)$$

where $\rho = -p$, which corresponds to the effective cosmological constant. Then if there is more than one solution satisfying $f'(q) = 0$ as $q = q_n$, $n = 0, 1, 2, \dots$, the theory could have effectively several different cosmological constants:

$$H = f(q_n), \quad \Lambda_n = 3f(q_n)^2. \quad (8)$$

Note that the solutions (8) corresponding to multiple cosmological constants may exist at the same moment which shows some analogy with cosmological landscape. Let us assume there is a solution corresponding to q_n . By perturbation, the solution may transit to another solution, say q_{n+1} . Hence, the transition period is proportional to $T_{n,n+1} = q_{n+1} - q_n$.

As an example, we consider the ideal fluid with the following time-dependent EoS

$$f(q) = f_0 - f_1 e^{-\lambda q} \left(\frac{\cos(\omega q + \alpha)}{\sqrt{\lambda^2 + \omega^2}} - \frac{1}{\lambda} \right). \quad (9)$$

Here f_0 and f_1 are constants and $f_0 > f_1 > 0$ and α ($0 < \alpha < \pi$, $\alpha \neq \pi/2$) is defined by

$$\tan \alpha = \frac{\omega}{\lambda}, \quad (10)$$

that is

$$\cos(\omega q + \alpha) = \frac{\lambda \cos \omega q - \omega \sin \omega q}{\sqrt{\lambda^2 + \omega^2}}. \quad (11)$$

Then one finds

$$f'(q) = f_1 e^{-\lambda q} (\cos \omega q - 1). \quad (12)$$

Since $f'(q) = 0$ when $q = \frac{2\pi n}{\omega}$ with an integer n , we have effectively multiple cosmological constant:

$$\Lambda_n = 3 \left\{ f_0 - f_1 e^{-2\pi n \lambda / \omega} \left(\frac{\cos(\alpha)}{\sqrt{\lambda^2 + \omega^2}} - \frac{1}{\lambda} \right) \right\}^2 = 3 \left\{ f_0 + \frac{f_1 e^{-2\pi n \lambda / \omega} \omega^2}{\lambda (\lambda^2 + \omega^2)} \right\}^2. \quad (13)$$

Eq.(5) shows also that there is a time dependent solution:

$$H(t) = f(t) = f_0 - f_1 e^{-\lambda t} \left(\frac{\cos(\omega t + \alpha)}{\sqrt{\lambda^2 + \omega^2}} - \frac{1}{\lambda} \right). \quad (14)$$

When $t = 2\pi n / \omega$, it follows $\rho = -p$ corresponding to the cosmological constants in (13). Note that the solutions with cosmological constant (13) may exist at the same moment. Such the universe could be considered as kind of multiverse [13] as it could be that some its regions may be inaccessible like in the model [14]. If we have a solution corresponding to q_n , by perturbation, the solution could transit to another solution corresponding to q_{n+1} . The time for transition could be given by $T = q_{n+1} - q_n = 2\pi / \omega$. Then the solution (14) describes the transitions between different Λ cosmologies (for related proposal of cascading universe, see [15]). Since $\Lambda_n > \Lambda_{n+1}$ in (13), the transition could occur from larger one to smaller one. In the limit of $t \rightarrow \infty$, $H(t)$ goes to f_0 , which corresponds to

$$\Lambda_\infty \equiv \lim_{n \rightarrow +\infty} \Lambda_n = 3f_0^2. \quad (15)$$

If $t \rightarrow \infty$ corresponds to the present time, we find $f_0 \sim 10^{-33}$ eV.

As a second slightly different example, one may consider the following ideal fluid:

$$f(q) = g_0 e^{-g_1 \left(q - \frac{1}{\omega} \sin \omega q \right)}, \quad (16)$$

which gives

$$f'(q) = -g_0 g_1 (1 - \cos \omega q) e^{-g_1 \left(q - \frac{1}{\omega} \sin \omega q \right)}. \quad (17)$$

In (16), it is assumed g_0 , g_1 , and ω are constants. Therefore, $f'(q) = 0$ when $q = \frac{2\pi n}{\omega}$ with an integer n , again. The effective multiple cosmological constant occurs:

$$\Lambda_n = 3g_0^2 e^{-\frac{4\pi n g_1}{\omega}}, \quad (18)$$

Eq.(16) shows that there exists also a time dependent solution:

$$H(t) = f(t) = g_0 e^{-g_1 \left(t - \frac{1}{\omega} \sin \omega t \right)}, \quad (19)$$

Again $t = 2\pi n / \omega$ corresponds to the cosmological constants in (16) and therefore the time-dependent solution could describe the transition between the cosmological constants, from larger one to smaller one. In the limit of $t \rightarrow +\infty$ or $n \rightarrow +\infty$, the effective cosmological constant vanishes: $\lim_{n \rightarrow +\infty} \Lambda_n = 0$, which is different from the example in (9) as found in (15).

It may be shown that the EoS (6) or (4) can be realized in the scalar-tensor theory, whose action is given by

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} R - \frac{1}{2} \omega(\phi) \partial_\mu \phi \partial^\mu \phi - V(\phi) \right\},$$

$$\omega(\phi) = -\frac{2}{\kappa^2} f'(\phi), \quad V(\phi) = \frac{1}{\kappa^2} (3f(\phi)^2 + f'(\phi)). \quad (20)$$

For the example (9), the corresponding scalar-tensor theory looks like

$$\omega(\phi) = -\frac{2f_1}{\kappa^2} e^{-\lambda \phi} (\cos \omega \phi - 1),$$

$$V(\phi) = \frac{1}{\kappa^2} \left\{ 3 \left(f_0 - f_1 e^{-\lambda \phi} \left(\frac{\cos(\omega \phi + \alpha)}{\sqrt{\lambda^2 + \omega^2}} - \frac{1}{\lambda} \right) \right)^2 + f_1 e^{-\lambda \phi} (\cos \omega \phi - 1) \right\}, \quad (21)$$

and for the example (16), it follows

$$\begin{aligned}\omega(\phi) &= \frac{2g_0g_1}{\kappa^2} (1 - \cos \omega\phi) e^{-g_1(1 - \frac{1}{\omega} \sin \omega\phi)} , \\ V(\phi) &= \frac{1}{\kappa^2} \left\{ 3g_0^2 e^{-2g_1(\phi - \frac{1}{\omega} \sin \omega\phi)} - g_0g_1 (1 - \cos \omega\phi) e^{-g_1(1 - \frac{1}{\omega} \sin \omega\phi)} \right\} .\end{aligned}\quad (22)$$

The appearance of exponential potentials may indicate to some connection with string theory. For other examples of scalar-tensor theory with de Sitter solutions, see [16].

Note that the solution (5) is always also a solution of the corresponding scalar-tensor theory (20) but the solution (7) is not always a solution of the scalar-tensor theory. This is because the fact that energy conservation law $\dot{\rho} + 3H(\rho + p) = 0$ is the first order differential equation while the scalar field equation is the second order differential equation:

$$0 = \omega(\phi)\ddot{\phi} + \frac{1}{2}\omega'(\phi)\dot{\phi}^2 + 3H\omega(\phi)\dot{\phi} + V'(\phi) . \quad (23)$$

Hence, even if the conservation law is satisfied, the field equation (23) is not always satisfied. The solution (7) requires that $\phi = q_0$ with $f'(q_0) = 0$ and therefore ϕ is a constant. Then in order that the solution (7) is also a solution of the scalar-tensor theory, we should require $V'(\phi) = 0$ at $\phi = q_0$. Since

$$V'(\phi) = \frac{1}{\kappa^2} \{ 6f(\phi)f'(\phi) + f''(\phi) \} , \quad (24)$$

if

$$f''(q_0) = 0 , \quad (25)$$

$\phi = q_0$ is also a solution of the scalar-tensor theory.

For the model (9) or (21), we find

$$f''(\phi) = f_1 e^{-\lambda\phi} \{ -\lambda(\cos \omega\phi - 1) - \omega \sin \omega\phi \} , \quad (26)$$

which vanishes when $\phi = \frac{2\pi n}{\omega}$ with an integer n . Since $f'(\phi)$ also vanishes there, $\phi = \frac{2\pi n}{\omega}$ is a solution of the scalar-tensor theory. Thus, the scalar-tensor theory has multiple cosmological constant solution

$$\frac{\Lambda_n}{\kappa^2} = V(2\pi n/\omega) = \frac{3}{\kappa^2} f(2\pi n/\omega)^2 , \quad (27)$$

which corresponds to (8). The first FRW equation (1) shows that

$$R = 12H^2 = 4\kappa^2\rho = 4\kappa^2V(2\pi n/\omega) = 4\Lambda_n , \quad (28)$$

and therefore the on-shell Lagrangian density $\mathcal{L}$ (20) is given by

$$\mathcal{L} = \frac{R}{2\kappa^2} - V(2\pi n/\omega) = \frac{\Lambda_n}{\kappa^2} = \frac{3}{\kappa^2} \left\{ f_0 + \frac{f_1 e^{-2\pi n\lambda/\omega\omega^2}}{\lambda(\lambda^2 + \omega^2)} \right\}^2 . \quad (29)$$

Hence, the energy of the corresponding cosmological constant solution changes with n , which is different from string landscape where millions of de Sitter vacua may have the same energy.

For the model (16) or (22), we find

$$f''(\phi) = -g_0g_1 \left\{ \omega \sin \omega\phi - g_1 (1 - \cos \omega\phi)^2 \right\} e^{-g_1(1 - \frac{1}{\omega} \sin \omega\phi)} . \quad (30)$$

Since both of $f'(\phi)$ and $f''(\phi)$ vanish when $\phi = \frac{2\pi n}{\omega}$ with an integer n , $\phi = \frac{2\pi n}{\omega}$ is also a solution of the corresponding scalar-tensor theory. Then the scalar-tensor theory has multiple cosmological constant solution (18). In this example, the corresponding on-shell Lagrangian density is given by

$$\mathcal{L} = \frac{3g_0^2}{\kappa^2} e^{-\frac{4\pi n g_1}{\omega}} . \quad (31)$$

3. Generalizing above EoS, we consider the case that $f(q)$ is given by two functions g and h in the following form:

$$f(q) = g(h(q) - \sin h(q)) . \quad (32)$$

Since

$$\begin{aligned} f'(q) &= g'(h(q) - \sin h(q)) (1 - \cos h(q)) h'(q) , \\ f''(q) &= g''(h(q) - \sin h(q)) (1 - \cos h(q))^2 (h'(q))^2 + g'(h(q) - \sin h(q)) (1 - \cos h(q)) h''(q) \\ &\quad + g'(h(q) - \sin h(q)) \sin h(q) (h'(q))^2 , \end{aligned} \quad (33)$$

one finds $f'(q) = f''(q) = 0$ when $h(q) = 2\pi n$ (n : integer) if g' , g'' , h' , and h'' are not singular. Hence, EoS (4) and the corresponding scalar-tensor theory (20) contains the multiple de Sitter solution which corresponds to the cosmological constant (18) when $h(q) = 2\pi n$ (n : integer):

$$\Lambda_n = 3 (g(2\pi n))^2 . \quad (34)$$

Moreover, such ideal fluid as well as corresponding scalar-tensor theory also leads to the time-dependent solution, whose Hubble rate is given by

$$H(t) = g(h(t) - \sin h(t)) . \quad (35)$$

Such (oscillating) universe passes through the de Sitter eras specified by different cosmological constants.

For instance, with the choice

$$h(q) = 2\pi N \tanh \omega q , \quad (36)$$

we have $2N - 1$ classical deSitter vacua, where $f'(q) = f''(q) = 0$ or $h(q) = 2\pi n$ with integers n 's ($-N < n < N$), between $-\infty < q < \infty$ and the limits $q \rightarrow \pm\infty$ also correspond to de Sitter vacua.

The effective EoS parameter w_{eff} is defined by

$$w_{\text{eff}} = -1 - \frac{2\dot{H}}{3H^2} . \quad (37)$$

When $w_{\text{eff}} < -1/3$, the universe is accelerating but when $w_{\text{eff}} > -1/3$, the universe is decelerating. As an example of (32), one can consider

$$f(q) = \frac{H_0}{H_1 + h(q) - \sin h(q)} , \quad (38)$$

where $h(q)$ is chosen as (36). Assume $H_1 > 2\pi N$ then $f(q)$ is not singular everywhere. One can now investigate the solution (35). Let define q_1 by

$$-2\pi(N-1) = h(q_1) = 2\pi N \tanh \omega q_1 . \quad (39)$$

Since $\dot{H} = 0$ when $t = -\infty$ or $t = q_1$, $w_{\text{eff}} = -1$ when $t = -\infty$ or $t = q_1$, that is, the universe is tentatively de Sitter space and therefore, the universe is accelerating. Hence, there should be an extremum of w_{eff} when $-\infty < t < q_1$. For the example (38), one gets

$$\dot{w}_{\text{eff}} = -\frac{2}{3H_0} \left\{ (1 - \cos h(t)) h''(t) + \sin h(t) (h'(t))^2 \right\} . \quad (40)$$

At the extremum, where $\dot{w}_{\text{eff}} = 0$, it follows

$$1 - \cos h(t) = -\frac{h'(t)}{h''(t)} \sin h(t) . \quad (41)$$

Let denote t satisfying the condition (41) by t_e , ($-\infty < t_e < t_1$). Note that t_e does not depend on H_0 . When $t = t_0$, it follows

$$w_{\text{eff}} = -1 + \frac{2(1 - \cosh h(t_e)) h'(t_e)}{H_0} = -1 + \frac{4\pi N\omega(1 - \cosh h(t_e))}{H_0 \cosh^2 \omega t_e} . \quad (42)$$

In (42), the second term is positive if $N\omega/H_0 > 0$. Since t_e does not depend on H_0 , we may choose H_0 to be small enough. Then w_{eff} can be larger than $-1/3$, that is, the universe can be decelerating. Since the universe is accelerating when $t = -\infty$ or $t = t_1$, the primordial universe, where $t \rightarrow -\infty$, is accelerating but when $t \rightarrow t_e$, the universe decelerates, that is, there occurs the transition from the accelerating universe to the decelerating universe. After that, when $t \rightarrow t_1$, the universe turns to acceleration again, that is, there occurs the transition from deceleration to acceleration. After t goes through t_1 , the universe passes through several de Sitter vacua and in the limit of $t \rightarrow +\infty$, the universe tends to the final de Sitter space where $H \rightarrow H_{\text{final}} = H_0/(H_1 + 2\pi N)$. $H(t)$ is monotonically decreasing function of t , the value of $H = H_{\text{final}}$ is minimum. The first period of the deceleration between $-\infty < t < t_1$ could be the order of $1/\omega$. Then in order to obtain the realistic model, we should choose ω to be the order of the age of the universe, that is, ten billion years or $\omega \sim 10^{-33} \text{ eV}$. When $t = t_1$, one has $H(t_1) = H_0/(H_1 - 2\pi(N-1))$. Since $H_1 > 2\pi N$, we find $H_1 - 2\pi(N-1) \sim \mathcal{O}(1)$ and the magnitude of $H(t_1)$ is the order of H_0 and therefore $H_0 \sim 10^{-33} \text{ eV}$, which corresponds to the observed value of the Hubble rate in the present universe. Since $H(-\infty) = H_0/(H_1 - 2\pi N)$, if the Hubble rate in the inflationary era is large enough, it follows $H_1 \sim 2\pi N$. Hence, the ideal fluid with time-dependent EoS of special form motivated by string landscape considerations may suggest the solution of the cosmological constant problem. Of course, similar scenario may be extended for scalar-tensor theory with corresponding potentials where also the unification of cosmological epochs may be achieved (for one related example, see [17]).

4. In the previous section, dark fluid describing the transitions between de Sitter eras is discussed. In this section, we consider if the transition between de Sitter and Anti-de Sitter space is possible in the similar framework. When cosmological constant is negative and there is no matter, there is no time evolution of the universe. One may include the matter with constant EoS parameter w , whose energy and the pressure are given by

$$\rho_m = \rho_0 a^{-3(1+w)} , \quad p_m = w\rho_0 a^{-3(1+w)} . \quad (43)$$

Then instead of (1), we find

$$\frac{3}{\kappa^2} H^2 = \rho + \rho_m , \quad -\frac{2}{\kappa^2} \dot{H} = p + \rho + p_m + \rho_m . \quad (44)$$

Hence, if ρ and p are given in terms of the function $g(q)$ with a parameter q as

$$\rho = \frac{3}{\kappa^2} g'(q)^2 - \rho_0 a_0^{-3(1+w)} e^{-3(1+w)g(q)} , \quad p = -\frac{3}{\kappa^2} f(q)^2 - \frac{2}{\kappa^2} f'(q) - w\rho_0 a_0^{-3(1+w)} e^{-3(1+w)g(q)} , \quad (45)$$

the following solution of (44) exists:

$$H = g'(t) \left(a(t) = a_0 e^{g(t)} \right) . \quad (46)$$

Note that the corresponding scalar theory is given by, instead of (21),

$$\begin{aligned} \omega(\phi) &= -\frac{2f_1}{\kappa^2} e^{-\lambda\phi} (\cos \omega\phi - 1) - \frac{w+1}{2} g_0 e^{-3(1+w)g(\phi)} , \\ V(\phi) &= \frac{1}{\kappa^2} \left\{ 3 \left(f_0 - f_1 e^{-\lambda\phi} \left(\frac{\cos(\omega\phi + \alpha)}{\sqrt{\lambda^2 + \omega^2}} - \frac{1}{\lambda} \right) \right)^2 + f_1 e^{-\lambda\phi} (\cos \omega\phi - 1) \right\} \\ &\quad + \frac{w-1}{2} g_0 e^{-3(1+w)g(\phi)} . \end{aligned} \quad (47)$$

Here

$$a_0 = \left(\frac{\rho_0}{g_0} \right)^{\frac{1}{3(1+w)}} . \quad (48)$$

We now consider the following model:

$$g(q) = \alpha(q) + \frac{2}{3(1+w)} \ln \sin \Omega q . \quad (49)$$

Here Ω is a constant and $\alpha(q)$ is a proper function which satisfies the conditions

$$\alpha(0) = \alpha(\pi/\Omega) = \alpha'(0) = \alpha'(\pi/\Omega) = 0 . \quad (50)$$

Assume, when $q \sim 0$,

$$\alpha(q) \sim \alpha_0 \Omega^2 q^2 , \quad (51)$$

and when $q \sim \pi/\Omega$,

$$\alpha(q) \sim \tilde{\alpha}_0 (\pi - \Omega q)^2 . \quad (52)$$

One gets

$$\rho = \frac{3}{\kappa^2} H^2 - \rho_m = \frac{3}{\kappa^2} \left(\alpha'(t) + \frac{2\Omega}{3(1+w)} \cot \Omega t \right)^2 - \rho_0 a_0^{-3(1+w)} e^{-3(1+w)\alpha(t)} \sin^{-2} \Omega t . \quad (53)$$

We now choose the parameters to satisfy the following condition:

$$\rho_0 a_0^{-3(1+w)} = \frac{4\Omega^2}{3(1+w)^2 \kappa^2} . \quad (54)$$

Then in the limit $t \rightarrow 0$, ρ goes to some constant:

$$\rho \rightarrow \frac{12\Omega^2}{\kappa^2(1+w)} \left(\alpha_0 - \frac{1}{9(1+w)} \right) , \quad (55)$$

and also in the limit $t \rightarrow \pi/\Omega$, ρ goes to the constant:

$$\rho \rightarrow -\frac{4\Omega^2}{\kappa^2(1+w)} \left(\tilde{\alpha}_0 + \frac{1}{3(1+w)} \right) . \quad (56)$$

Hence, in the limits of $t \rightarrow 0$, π/Ω , ρ behaves as a cosmological constant. For example, if it is chosen

$$\alpha_0 > \frac{1}{9(1+w)} , \quad \tilde{\alpha}_0 > -\frac{1}{3(1+w)} , \quad (57)$$

ρ behaves as a positive cosmological constant when $t \sim 0$ and as a negative cosmological constant when $t \sim \pi/\Omega$. Therefore, the dark fluid (49) could describe the transition between universe regions with positive and negative cosmological constants.

5. In summary, we discussed FRW universe with time-dependent EoS dark fluid as a toy model which leads to multiple de Sitter epochs. This suggests some classical analog of cosmological landscape. It remains to show which stringy compactification may be described as an effective ideal fluid of such (or similar) form.

The possibility of transition between different de Sitter eras is demonstrated in the universe expansion history. That may suggest the reasonable approach to the solution of cosmological constant problem, explaining

its effective decrease. In addition, the possibility of transition between regions with positive and negative cosmological constant is demonstrated. It is not difficult to generalize the above study for the presence of standard CDM matter. In such a way, the standard Λ CDM cosmology which is known to be consistent with observational data (for recent review, see [18]) easily emerges with reasonable explanation of the smallness of current cosmological constant.

Acknowledgements. The work by S.N. was supported in part by Monbusho grant no.18549001 (Japan) and 21st Century COE program of Nagoya Univ. provided by JSPS (15COEEG01) and that by S.D.O. by the project FIS2005-01181 (MEC, Spain), by AGAUR (Generalitat de Catalunya), contract 2005SGR-00790, by LRSS project N4489.2006.02 and by RFBR grant 06-01-00609 (Russia).

-
- [1] E. Copeland, M. Sami and S. Tsujikawa, arXiv:hep-th/0603057; T. Padmanabhan, Phys. Repts. **380** (2003) 235.
 - [2] V. Cardone, C. Tortora, A. Troisi and S. Capozziello, Phys. Rev. D **73** (2006) 043508.
 - [3] S. Nojiri and S. D. Odintsov, Phys. Rev. D **70** (2004) 103522 [arXiv:hep-th/0408170]; H. Stefancic, Phys. Rev. D **71** (2005) 084024.
 - [4] S. Nojiri and S. D. Odintsov, Phys. Rev. D **72** (2005) 023003 [arXiv:hep-th/0505215]; Phys. Lett. B **639** (2006) 144 [arXiv:hep-th/0606025]; S. Capozziello, V. Cardone, E. Elizalde, S. Nojiri and S. D. Odintsov, Phys. Rev. D **73** (2006) 043512 [arXiv:astro-ph/0508350].
 - [5] I. Brevik and O. Gorbunova, GRG **37** (2005) 2039 [arXiv:gr-qc/0504001]; J. Ren and X. Meng, Phys. Lett. B **633** (2006) 1 [arXiv:astro-ph/0511163]; I. Brevik, arXiv:gr-qc/0601100.
 - [6] S. Capozziello, S. Nojiri and S. D. Odintsov, Phys. Lett. B **634** (2006) 93 [arXiv:hep-th/0512118]; S. Capozziello, S. Nojiri, S. D. Odintsov and A. Troisi, Phys. Lett. B **639** (2006) 135 [arXiv:astro-ph/0604431].
 - [7] S. Nojiri and S. D. Odintsov, Phys. Rev. D **68** (2003) 123512, arXiv:hep-th/0307288; arXiv:hep-th/0601213.
 - [8] I. Arefeva, L. Joukovskaya and S. Vernov, arXiv:hep-th/0701184; A. Koshelev, arXiv:hep-th/0701103.
 - [9] L. Susskind, arXiv:hep-th/0302219; M. Douglas, JHEP **0305** (2003) 046 [arXiv:hep-th/0303194]; T. Banks, M. Dine and E. Gorbatov, JHEP **0408** (2004) 058 [arXiv:hep-th/0309170];
 - [10] M. Bastero-Gil, K. Freese and L. Mersini-Houghton, Phys. Rev. D **68** (2003) 123514.
 - [11] V. Kozlov and I. Volovich, arXiv:hep-th/0612135.
 - [12] S. Nojiri and S. D. Odintsov, GRG **38** (2006) 1285 [arXiv:hep-th/0506212]; S. Capozziello, S. Nojiri and S. D. Odintsov, Phys. Lett. B **632** (2006) 597 [arXiv:hep-th/0507182].
 - [13] S. Robles-Perez, P. Martin-Moruno, A. Rozas-Fernandez and P. Gonzalez-Diaz, arXiv:gr-qc/0701127.
 - [14] B. McInnes, arXiv:gr-qc/0611101.
 - [15] S. Watson, M. Perry, G. Kane and F. Adams, arXiv:hep-th/0610054.
 - [16] V. Faraoni, Phys. Rev. D **70** (2004) 044037; Z. Huang, H. Lu and W. Fang, Class. Quant. Grav. **23** (2006) 6215.
 - [17] S. Nojiri and S. D. Odintsov, Phys. Lett. B **562** (2003) 147 [arXiv:hep-th/0303117].
 - [18] L. Perivolaropoulos, arXiv:astro-ph/0601014.